



PRODUCED AND MANUFACTURED IN AUSTRALIA
 PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@pickrwl.com.au • www.spraygro.com.au
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 499
 52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE
SPRAYGRO LIQUID FERTILIZERS

Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.
STORAGE & DISPOSAL:

ANALYSIS	AUSTRALIA W/V %	INTERNATIONAL W/V %
OLIVE TREES	15-18	3-5
PAW PAW	8-10	1-3
PEANUTS	20-25	N/A
PINEAPPLES	6-10	1-3
POTATOES	5-10	2-4
RED WINE	10-20	3-5
GRAPES	5-7	N/A
WHITE WINE	8-15	2-3
GRAPES	8-15	2-3
TOMATOES	7-10	N/A
TABLE GRAPES	10-15	N/A
TURF	N/A	300-500ml/100m ²
Chelated Zinc	1	1
Potassium K	10	10
Phosphorus P	20	30 P ₂ O ₅
Nitrogen N	18	12
Specific gravity	1.50	1.07

CROP	Fertilization L/ha	Foliar L/ha
CANOLA	20-30	N/A
LEGUMES	20-25	N/A
MAIZE/ SUGARCANE	15-20	N/A
LUCERNE	N/A	3-5
BANANAS	8-12	N/A
BEANS	4-6	N/A
BRASSICAS	6-10	2-3
CUCURBITS	5-10	2-3
LEAFY GREENS	7-10	N/A

BROADACRE: Foliar rate of application per min water/hectare: 50 litres.
HORTICULTURE: Foliar rate of application per min water/hectare: 150-350 litres. Tree crops may require higher rates (500-1000L).
 Apply through soil injection at planting. Apply through soil injection at planting. Reduce rates 5-6L/ha) with Anhydrous ammonia, if has been injected. Fertigation again at early flowering.
 Apply at planting in conjunction with 42N.
 Apply every ration crop.
 Apply when sufficient leaf area exists after every cut.
 Apply to juvenile trees at early establishment-ferigation. Apply foliar at least two weeks before flowering.
 Apply early in the season. Repeat if necessary. Apply through soil after transplant. Apply foliar at early heading.
 Repeat if necessary.
 Apply every 14 days from mid-crop through to harvest.



LIQUID FERTILIZER

CHARGE

N P K
18 : 20 : 10 w/w
+ 1% Zn w/w

20 Litre NET 200 Litre NET 1000 Litre NET

CAUTION

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

Safety glasses, gloves and protective clothing should be worn when using this product.

DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred of suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.



BATCH No.