

**APPLICATION RECOMMENDATIONS: SPRAYGRO FIRMRITE:**

**HORTICULTURE:**

**Rate of application per min water/hectare:** 300 litres

HORTICULTURE CROPS @ 5-10 litres/hectare. Apply as needed or from when plants are 10-15cm in height and in rapid growth stage. Apply every 14-21 days.

CITRUS @ 10 litres/hectare. Apply 3 applications at bud burst 7-14 days apart.

NAVEL ORANGES @ 20 litres/hectare. To assist in reduction of albedo breakdown, apply 10 days apart from mid January to mid February.

VINES @ 5-10 litres/hectare. Apply at regular intervals from fruit formations.

LETTUCE @ 5-10 litres/hectare. Apply from emergence at 14 day intervals as required. Aids in reduction of tip burn.

**FRUIT CROPS:**

**Rate of application per min water/hectare:** 350 litres

APPLES @ 7-10 litres/hectare. Apply at petal fall. Continue at 7-14 days apart until harvest. Aids in reduction of Bitter Pit.

POST HARVEST/DIP @ 600mls per 100 litres. Other apple varieties at 2-3 litres per 100 litres.

GOLDEN DELICIOUS @ 2-4 litres/hectare. Apply as required to improve firmness and quality.

FRUIT, TROPICAL @ 2-5 litres/hectare. Apply at 10-14 day intervals during flowering.

STRAWBERRIES @ 7-16 litres/hectare. Apply at flowering and as required, enhances seed set.

TOMATOES

**LIQUID FOLIAR FERTILIZER**

**FIRMRITE™**

HIGH CALCIUM SOLUTION DESIGNED TO INCREASE FRUIT DEVELOPMENT, FIRMNESS & MATURITY. CALCIUM STRENGTHENS CELL WALLS IN FRUIT & VEGETABLES.

**17% W/V CALCIUM**

**20 Litre NET**  **200 Litre NET**  **1000 Litre NET**



**CAUTION**

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- **Compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**Safety glasses, gloves and protective clothing should be worn when using this product.**

**CONDITIONS OF SALE:** 1) Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than the officer of the Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2) Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

**SPRAYGRO LIQUID FERTILIZERS PTY. LTD.**

52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE  
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496  
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@picknowl.com.au

**BATCH No.**

**PRODUCED AND MANUFACTURED IN AUSTRALIA**