



BATCH No.

**DISCLAIMER/CONDITIONS OF SALE:** (1) Foliar Fertilizer Pty Ltd trading as SprayGro Liquid Fertilizer and its servants and agents ("SprayGro") shall be under no liability to the purchaser and/or user of SprayGro products ("Customer") in respect of any loss or damage (including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the SprayGro Product Profile or not. (2) The customer warrants that there has been no reliance by the customer upon any representation made by SprayGro or upon and descriptions, illustrations or specifications contained in SprayGro's Product Profile or any other document produced by SprayGro. (3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. (4) The customer shall at all times indemnify and hold harmless SprayGro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred or suffered by SprayGro arising from any proceedings or threatened proceedings against SprayGro by or involving the customer. (5) SprayGro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. (6) No other than an authorised officer of SprayGro is authorised to make any warranty, guarantee or direction concerning this product. (7) Before using please obtain a current Product Profile which is available from SprayGro or our Distributor.

**Safety glasses, gloves and protective clothing should be worn when using this product.**

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

- DO NOT apply during the heat of the day.
- DO NOT spray to run off. Do not spray to visible droplet coalescence. Do not allow concentrated spray mist to run on fruit or leaves.
- DO NOT apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- DO NOT use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- Use the correct label spray rate of any pesticide used with SprayGro product.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product. do not proceed further.
- When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. damage to the plant.

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in

**CAUTION**

0141852

**LIQUID FERTILIZER**

**FIRMRITE™ CB**

HIGH CALCIUM / BORON SOLUTION DESIGNED TO INCREASE FRUIT SET DEVELOPMENT, FIRMNESS & MATURITY, CALCIUM STRENGTHENS CELL WALLS IN FRUIT & VEGETABLES.

**16% W/W CALCIUM**  
**0.2% W/W BORON**

20 Litre NET  200 Litre NET  1000 Litre NET

**HORTICULTURE:** Foliar applications per min water/hectare generally required: 300L. Tree crops may require higher rates (500-1000L).

CROP	FOLIAR RATES L/ha	COMMENTS
ASPARAGUS	2-4L	Apply 7 days from 1st fern. Repeat at spear emergence.
AVOCADOES	3-7L	Apply 3 weeks prior to flowering. Repeat every 3 weeks from fruit set to harvest.
BROCCOLI	2-4L	Apply every 2 weeks until head formation.
CAPSICUM	4-5L	Apply every 2 weeks until flowering. Repeat at regular intervals during fruiting.
CELERY	2-4L	Apply every 2 weeks. Stop application 1 week before harvest.
CITRUS	5-6L	Apply prior to bloom, at petal fall and later as required.
COTTON	5-6L	Apply prior to 1st flowering. Repeat at boll formation.
CUCUMBERS & CUCURBITS	2-5L	Apply prior to flowering with further applications at regular intervals. Do not apply a week before harvest.
GRAPE VINES	4-6L	Apply prior to flowering with regular applications until 2-3 weeks before harvesting. Minimum water 500L/ha.
LETTUCE	5-10L	Apply from emergence, continue 14 day intervals until harvest.
NUT CROPS	2-4L	Apply 2 weeks post bloom, repeat at regular intervals during nut development. Repeat post-harvest.

CROP	FOLIAR RATES L/ha	COMMENTS
OLIVES	3-4L	Apply 2 weeks prior to fruit set, at fruit set and further applications as required.
POME FRUIT	5-10L	Apply at petal fall. Continue every 14 days to harvest.
POTATOES	4-5L	Apply 3 weeks after emergence. Repeat at hooking, canopy closure and bulking.
STONE FRUITS	5-10L	Apply post-bloom. Continue from fruit set at regular intervals.
STRAWBERRIES	2-4L	Apply 2 applications prior to flowering with further applications at regular intervals.
TOMATOES	7-16L	Apply at flowering then at 14 day intervals.
TROPICAL FRUIT	2-4L	Spray before bloom. Repeat at fruit set until 7 days before harvest.
VEGETABLES	2-5L	Apply 3 weeks after emergence or post-transplant. Repeat at regular intervals.
ZUCCHINI	4-5L	Two applications prior to flowering. Repeat during fruit elongation until 1 week before harvest.

ANALYSIS	AUSTRALIA W/W %	INTERNATIONAL W/W %
Calcium (Ca)	16.0	12
Boron (B)	0.2	0.15
Specific gravity 1.33		



NOT APPLICABLE TO 1000 LITRE CONTAINERS

**STORAGE & DISPOSAL:**

Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

**SPRAYGRO LIQUID FERTILISERS**

52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE PORT ADELAIDE, SOUTH AUSTRALIA 5015  
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • E-mail: spraygro@spraygro.com.au • www.spraygro.com.au • A.B.N. 47 007 974 496

**PRODUCED AND MANUFACTURED IN AUSTRALIA**