



## LIQUID FERTILIZER

# MAXI-PHOS INJECTA-23

N P K  
16 - 23 - 0 W/W

Nitrogen as ammonia .....16% W/W  
Phosphorous as Poly Phosphate.....23% W/W

Can be used as a superior liquid alternative to granular MAP. More available form of Phosphorous.

20 Litre NET  200 Litre NET  1000 Litre NET

Maxi-Phos Injecta-23 application recommendations: Apply to soil via ground injection, band spraying or via fertigation. Primarily used as a starter fertilizer or when phosphorous availability to crops is limited. Apply based on amount of P required.

CROP	FERTIGATION RATES L/ha
ALMONDS	20-30L
APPLES	10-20L
ASPARAGUS	10-15L
AVOCADOS	10-15L
BANANAS	10-20L
BROCCOLI	10-20L
CAPSICUM	10-25L
CITRUS	30-40L
CUCUMBERS	10-15L
GRAPE VINES	20-40L
LETTUCE	10-15L

CROP	FERTIGATION RATES L/ha
MELONS	15-25L
NUT CROPS	15-20L
OLIVES	15-30L
POME FRUIT	15-25L
POTATOES	20-40L
STONE FRUIT	15-25L
STRAWBERRIES	15-20L
TOMATOES	15-25L
TROPICAL FRUIT	10-20L
VEGETABLES	10-15L
ZUCCHINI	10-15L

**BROADACRE:**  
RATE OF SOIL APPLICATION PER MIN WATER/HECTARE: 100 litres.  
CEREALS, COTTON: Apply Maxi-Phos @ 26-45 litres/hectare.

- Note: These rates provide 6-10kg P/ha.**
- BRASSICAS, LEGUMES: Apply Maxi-Phos @ 23-41 litres/hectare.
  - Ideally, inject as a continuous band 2-3cm below the seed - avoid dribble out of nozzles as this will increase the chance of a root intercepting the fertilizer.
  - Maxi-Phos Injecta can also be applied into the soil surface by using high pressure nozzles, followed by press wheels.
  - SARDI research in 2001 found no conclusive differences between the performance of spray or stream nozzles.

**HORTICULTURE:**  
RATE OF APPLICATION PER MIN WATER/HECTARE: 1000 litres.  
FOLIAR @ 7-10 litres/hectare.  
FERTIGATION minimum water rate is 1:100.

**NOTE: Only mix enough Maxi-Phos Injecta for 1 days application. Do not store Maxi-Phos mixed with water or chemicals.**

ANALYSIS	AUSTRALIA W/W %	INTERNATIONAL W/W %
Nitrogen N	16	11
Phosphorous P	23	P.O-37
Potassium K	0	K.O.0

Specific gravity 1.42 pH 6.5-6.9



Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

**STORAGE & DISPOSAL:**

SPRAYGRO LIQUID FERTILIZERS  
52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE PORT ADELAIDE, SOUTH AUSTRALIA 5015  
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au • A.B.N. 47 007 974 498

**CAUTION**


The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture.
- Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**Safety glasses, gloves and protective clothing should be worn when using this product.**

**DISCLAIMER/CONDITIONS OF SALE:** 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.



BATCH No.