

BATCH NO.



**DISCLAIMER/CONDITIONS OF SALE:** (1) Foliar Fertilizers Pty Ltd trading as SprayGro Liquid Fertilizer and its servants and agents ("SprayGro") shall be under no liability to the purchaser and/or user of SprayGro products ("Customer") in respect of any loss or damage (including consequential loss or damage which may be suffered, incurred or arise directly or indirectly) in respect of such products, or the use thereof, in accordance with the SprayGro Product Profile or not. (2) The customer warrants that there has been no reliance by the customer upon any other representation made by SprayGro or upon and descriptions, illustrations or specifications contained in SprayGro's Product Profile or any other document produced by SprayGro. (3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. (4) The customer shall at all times indemnify and hold harmless SprayGro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred or suffered by SprayGro arising from any proceedings or proceedings threatened against SprayGro by or involving the customer. (5) SprayGro warrants that this product consists of ingredients specified and is responsible for the purpose stated on this label when used in accordance with the directions under normal conditions of use. No other than an authorised officer of SprayGro is authorised to make any warranty, guarantee or direction concerning this product. (6) Before using please obtain a current Product Profile which is available from SprayGro or our Distributor.

**Safety glasses, gloves and protective clothing should be worn when using this product.**

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be observed and observed prior to the total crop spray.

- **DO NOT** apply during the heat of the day.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- bucket (test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture.

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

**CAUTION**

44181N2



LIQUID FERTILIZER

RESULT<sup>®</sup>

PLUS

N P K  
14 : 8 : 10 w/v

20 Litre NET ■ 200 Litre NET ■ 1000 Litre NET ■

Application recommendations for Result Plus 14-8-10. Apply with sufficient water for thorough coverage of leaves when foliar applying.

HORTICULTURE: Foliar rate of application per min water/hectare:

CROP	FOLIAR RATES L/ha	FERTIGATION RATES L/ha	COMMENTS
AVOCADOES AND TROPICAL FRUIT	4L	15L	Apply at 7-14 day intervals from fruit set through to harvest.
ALMONDS	2-3L	15-30L	Apply post-bud burst at 14 day intervals until hull split.
ASPARAGUS	2-4L	7-10L	Apply at spear emergence until harvest.
CITRUS	2-4L	15-30L	Apply post-flowering until fruit expansion.
CAPSICUM	3-4L	10-30L	Apply post-planting until mid fruiting.
BRASSICA'S	4L	10-15L	Apply at early head development to harvest every 14 days.
CUCURBITS	2-4L	15-20L	Apply at fruit set onwards every 7-14 days until harvest.
GARLIC AND ONIONS	3-4L	12-15L	Apply from when sufficient leaf area exists to intercept spray and at bulb development through to harvest.
GRAPES	3-5L	15-30L	Apply post-bud burst until veraison.
LEAFY GREENS	2-3L	8-12L	Apply every 7-14 days from 4 leaf stage until harvest.
OLIVE TREES	3-4L	15-20L	Apply from early through to late fruiting.

CROP	FOLIAR RATES L/ha	FERTIGATION RATES L/ha	COMMENTS
PAW PAW	2-3L	8-12L	Apply at end of flowering and 21 days after flowering as required.
PEANUTS	n/a	10-20L	Apply at planting.
PINEAPPLES	2-3L	5-10L	Apply at 14-21 day intervals from fruit set through to harvest.
POTATOES	4-5L	12-20L	Apply every 7-14 days from tuber initiation to harvest.
STRAWBERRIES	2-4L	10-20L	Apply post-planting at regular intervals.
TOMATOES	2-3L	10-15L	Apply at early to late vegetative growth.
VEGETABLES	4-6L	10-15L	Apply at regular intervals during growing season.

**BROADACRE: Foliar rate of application per min water/hectare:** 50 litres.  
 COTTON @ 6-9 litres/hectare at 4-6 leaf stage and during mid to late flowering.  
 CEREALS @ 6-9 litres/hectare when sufficient leaf area exists to spray.  
 LEGUME GRAINS @ 4 litres/hectare during pod development.  
 TURF @ 400-500mls/100m<sup>2</sup>. Water dilution 1:100.

ANALYSIS	AUSTRALIA W/V %	INTERNATIONAL W/V %
Nitrogen N	14	10.85
Phosphorus P	8	14.20 (P <sub>2</sub> O <sub>5</sub> )
Potassium K	10	9.30 (K <sub>2</sub> O)

+ edta Chelated trace elements Fe 0.03%, Mn 0.03%, Zn 0.02%, Cu 0.02%, Mg 0.02%, Mo 0.001%, B 0.017%, Co 0.001%, S 0.05%. Specific gravity 1.29

**STORAGE & DISPOSAL:**

Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

**SPRAYGRO LIQUID FERTILIZERS**

52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE PORT ADELAIDE, SOUTH AUSTRALIA 5015  
 PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au • A.B.N. 47 007 974 496



NOT APPLICABLE TO 1000 LITRE CONTAINERS

PRODUCED AND MANUFACTURED IN AUSTRALIA