



- APPLICATION RECOMMENDATIONS:** SPRAYGRO SMARTRACE ZINC-COPPER is suitable for the following crops where a zinc & copper deficiency exists. Apply with sufficient water for thorough coverage of leaves.
- BROADACRE:** Rate of application per min water/hectare: 50 litres
- ③ 3-4 litres/hectare. CEREAL
 - ③ 3-4 litres/hectare. LEGUMES, CANOLA
 - ③ 2-5 litres/hectare. PASTURE
 - ③ 2-3 litres/hectare. COTTON
 - ③ 2-4 litres/hectare. HORTICULTURE: Rate of application per min water/hectare: 300 litres.
 - ③ 2-4 litres/hectare. CITRUS, POME FRUIT,
 - ③ 2-4 litres/hectare. POPPES
 - ③ 3-4 litres/hectare. ALMONDS
 - ③ 1-2 litres/hectare. VINES
 - ③ 1-3 litres/hectare. STRAWBERRIES
 - ③ 2-4 litres/hectare. SILVICULTURE
 - ③ 2-4 litres/hectare. VEGETABLES
 - ③ 150ml/100m²/water dilution 1:100. TURF
- SEED DRESSINGS:** Rate per tonne:
- ③ 7-9 litres. CEREALS
 - ③ 8-10 litres. LEGUME GRAINS
 - ③ 10 litres/tonne of seed. MAZE, RICE
 - ③ 9-10 litres. CANOLA

The rates must be adjusted according to the tissue and/or soil analysis results.

ANALYSIS	AUSTRALIA W/V %	INTERNATIONAL W/V %
Zinc Zn	3.0	2.4
Copper Cu	6.0	4.8
Specific gravity	1.25	pH 3.0 - 4.0

Easily Applied. No dust residues. Requires no added water. Physically compatible with most pickles (known to be not compatible with P-Pickle™ and Jockey™).

Apply in springtime as a thorough spray, up until bloom if necessary. Repeat applications may be necessary for a severe deficiency. Best absorption occurs 1/2 - 1/3 through a flush of growth. Spray should not be applied during blossoming or harvest.

Apply when trees are completely dormant (between late May and before budswell begins). Do not apply during bloom. Repeat as necessary. Apply a week after transplanting then again prior to bloom if necessary. Young foliage, flowers or fruit are susceptible to trace-element spray damage. Test a few plants or a plot before application.

Apply when deficiency is known to exist. Repeat as necessary. Apply when plants are young and before flowering. Repeat if necessary.

Apply during spring if a deficiency exists. Repeat as necessary.

SPRAYGRO LIQUID FERTILIZERS
52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au
PRODUCED AND MANUFACTURED IN AUSTRALIA

SPRAYGRO®

SMARTRACE®

CHELATED ELEMENT

ZINC-COPPER

CONTAINS
5.0% W/W
SULPHUR

GUARANTEED MINIMUM ANALYSIS

ZINC AS NATURAL PHENOLIC COMPLEX 6.0% W/W

COPPER AS NATURAL PHENOLIC COMPLEX 3.0% W/W

20 Litre NETO 200 Litre NETO 1000 Litre NETO

CAUTION

- The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.
- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
 - Physically compatible with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
 - Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
 - Use the correct label spray rates of any pesticide used with SprayGro product.
 - DO NOT use when crop is under heavy stress from pests, heat or inadequate moisture levels.
 - DO NOT apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
 - DO NOT spray to run off. Do not spray to visible droplet coalescence. Do not allow concentrated spray mist to run on fruit or leaves.
 - DO NOT apply during the heat of the day.

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

Safety glasses, gloves and protective clothing should be worn when using this product.

DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred of suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.



BATCH No.