


BATCH No.



Safety glasses, gloves and protective clothing should be worn when using this product.

and observed prior to the total crop spray.

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• Use the correct label spray rates of any pesticide used with SprayGro product.

• Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

LIQUID FERTILIZER

SECURE AN-28

N : P : K
28 : 0 : 0 W/W

NITROGEN AS AMMONIUM 14.0% W/W.
NITROGEN AS NITRATE 14.0% W/W.

20 Litre NET ■ 200 Litre NET ■ 1000 Litre NET ■

Secure AN-28 application recommendations (for Secure AN-25 adjust rates accordingly to analysis variation):
Minimum water rate is 150 for foliar application and 1:100 for fertigation. Normal application rates should be based on the amount of N required.

HORTICULTURE: Rate of foliar application per min water/hectare: 1000 litres.

CROP	FOLIAR RATES L/ha	FERTIGATION RATES L/ha	COMMENTS
ALMONDS	4-6L	15-20L	Apply at regular intervals post flowering.
APPLES	5-8L	10-20L	Apply in spring and summer at regular intervals.
ASPARAGUS	4-6L	7-10L	Apply at planting, first year and at regular intervals.
AVOCADOS	6-9L	7-10L	Apply to meet N requirement.
BANANAS	6-9L	8-10L	Apply as and when required to stimulate growth.
BROCCOLI	5-8L	7-10L	Apply post emergence at regular intervals according to N demand.
CAPSICUM/PEPPERS	3-4L	10-20L	Apply post emergence at regular intervals.
CITRUS	6-12L	15-20L	Apply at regular intervals according to N demand.
CUCUMBERS	3-4L	10-15L	Monthly applications commencing 4 weeks post emergence or 21 days after transplanting.
GRAPE VINES	4-6L	10-15L	Apply every 14 days during the growing season and once post harvest.
LETTUCE	4-6L	10-15L	Apply post emergence/transplanting and at regular intervals.
MELONS	4-8L	12-20L	Apply post emergence/transplanting and at regular intervals.
NUT CROPS	5-9L	15-20L	Apply at regular intervals post flowering and once post harvest.
OLIVES	6-10L	15-20L	Apply in spring and during growth phase at regular intervals: apply post harvest.

BROADACRE: Rate of foliar application per min water/hectare: 80 litres.

CROP	FOLIAR RATES L/ha	FERTIGATION RATES L/ha	COMMENTS
POTATOES	6-12L	10-20L	Apply monthly 2-3 weeks after emergence.
STONE/POPE FRUIT	6-12L	12-24L	Apply in spring and summer after stem extension growth.
STRAWBERRIES	5-8L	15-20L	Apply post emergence at regular intervals according to N demand.
TOMATOES	6-9L	10-20L	Apply post emergence at regular intervals according to N demand.
VEGETABLES	5-8L	7-10L	Apply monthly commencing post emergence or 21 days after transplanting.
ZUCCHINI	6-8L	9-12L	Apply monthly commencing post emergence or 21 days after transplanting.

TURF: Rate of application 6-800ml/100m²water dilution 1:100.

ANALYSIS	AUSTRALIA W/W %	INTERNATIONAL W/W %
Nitrogen N	28	22.3
Phosphorus P	0	P:O:0
Potassium K	0	K:O:0

Specific gravity 1.25 pH 5-6 Salting out temperature 5°C

NOTE: SECURE AN-28 has a salting out temperature of less than 5°C (SECURE AN-25 is around 2°C). Product will easily redissolve as soon as it is exposed to warmer temperatures again. Otherwise add warm water to product.

MATERIAL OF CONSTRUCTION: SECURE AN-28/25 are highly corrosive. Aluminium, stainless steel and certain fiberglass products are good for storage and applicator tanks.

STORAGE & DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

SPRAYGRO LIQUID FERTILISERS

52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 FORT ADELAIDE BUSINESS CENTRE PORT ADELAIDE, SOUTH AUSTRALIA 5015
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au • A.B.N. 47 007 974 496



NOT APPLICABLE TO 1000 LITRE CONTAINERS

PRODUCED AND MANUFACTURED IN AUSTRALIA