



**PRODUCED AND MANUFACTURED IN AUSTRALIA**

**SPRAYGRO LIQUID FERTILIZERS**  
 52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE  
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496  
 PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@pickknow.com.au • www.spraygro.com.au

**STORAGE & DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

ANALYSIS	AUSTRALIA W/V %	INTERNATIONAL W/V %
Nitrogen N	12	8.6
Phosphorus P	4	6.45 P <sub>2</sub> O <sub>5</sub>
Potassium K	23	20.1 K <sub>2</sub> O
Humates	3	2.15

**NB: DO NOT APPLY AS FOLIAR AS IT MAY STAIN FRUITS.**

CROP	Fertigation L/ha	Water Dilution 1:60-100
RED WINE GRAPES	12-16	Apply at vine establishment. Repeat 14-21 day intervals. Fertigate post-harvest.
STRAWBERRIES	7-10	Apply at planting. Repeat 14-21 days, if required. Do not apply at peak fruiting.
TOMATOES	8-12	Apply at transplant. Repeat at truss formation. Do not apply at peak fruiting.
CROP	5-7	Apply via water injection or furrow spray at planting. Do not apply on foliage.
COTTON	5-7	Apply after each cut.
LUCERNE	7-10	Apply at early germination or at green construction/renovation.
TURF	350-500ml/100m <sup>2</sup>	

CROP	Fertigation L/ha	Water Dilution 1:60-100
AVOCADOES	15-18	Apply to juvenile trees at early establishment. Repeat as required. Do not apply on foliage.
TROPICAL FRUIT	10-15	Apply early season. Repeat at fruiting. Do not apply on foliage.
BANANA	8-10	Apply at planting. Repeat, if required, at fruiting. Do not apply to foliage.
BEANS	5-10	Apply at planting. Repeat at fruit set.
BRASSICAS	8-10	Apply after transplanting. Repeat at heading.
CUCURBITS	7-10	Apply at 1-2 leaf stage. Repeat at 4-5 leaf stage. Apply before fruit set.
GARLIC/ONIONS	10-12	Apply 1 week after emergence. Repeat 14-21 day intervals.
LEAFY GREENS	12-15	Apply at planting/3 days after transplanting. Repeat 14-21 day intervals.
OLIVE TREES	15-20	Apply early season. Repeat at peak fruiting. Apply post harvest to maintain tree vigour.
PAW PAW	10-15	Apply early season. Repeat at flowering and fruiting stages.
PINEAPPLES	8-12	Apply early season. Repeat as required.
POTATOES	10-15	Apply at planting. Repeat as required. Do not apply at tuberization.



**LIQUID FERTILIZER**

**TOWER**

**N P K**  
**12 : 4 : 23 W/V**

**20 Litre NET • 200 Litre NET • 1000 Litre NET**

**CAUTION**

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**Safety glasses, gloves and protective clothing should be worn when using this product.**

**DISCLAIMER/CONDITIONS OF SALE:** 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred of suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.



BATCH No.