


BATCH No.



**DISCLAIMER/CONDITIONS OF SALE:** 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.

**Safety glasses, gloves and protective clothing should be worn when using this product.**

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

- Use the correct label spray rates of any pesticide used with SprayGro product. Surfactants are not recommended for use with Zintron since they may increase phytotoxicity.

- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further. Do not mix with phosphate fertilizers.

- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.

- Use the correct label spray rates of any pesticide used with SprayGro product. Surfactants are not recommended for use with Zintron since they may increase phytotoxicity.

- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

- **DO NOT** apply during the heat of the day.



20 Litre NET  200 Litre NET  1000 Litre NET

CONTAINS Zn AS ZINC NITRATE FOR MAXIMUM FOLIAR UPTAKE.

# ZINTRON

## 19-0-0 + 6.6Zn

LIQUID FERTILIZER

Zintron application recommendations: Apply when zinc deficiency exists. Horticulture - Foliar applications per min water/hectare spray to cover generally. 150-300L. Tree crops may require higher rates 500-1000L.

CROP	FOLIAR RATES L/ha	COMMENTS
ALMONDS	1-3L	Apply at petal fall, repeat every 4 weeks, post harvest.
APPLES	1-3L	Apply at bud break through full bloom, post harvest.
APRICOTS	0.5-1L	Apply post harvest, dormant only.
ASPARAGUS	1-4L	Apply at full fern. One application per year.
AVOCADO	1-3L	Best applied at Spring flush. Repeat as required.
BEANS	1-4L	Apply at sufficient foliage, repeat 100 up to full bloom.
BLUEBERRIES	0.5-1L	Apply from leaf expansion up to small fruits.
BROCCOLI	1-4L	Apply first at 150-200mm. Repeat 7-10 days.
BRUSSELS SPROUTS	1-2L	Apply first at 150-200mm. Repeat 7-10 days.
CABBAGE	1-4L	Apply at 100 to 2 weeks post thinning. Repeat if required.
CARROTS	1-4L	Apply at 100-150mm tall. Repeat in 10-14 days.
CELERY	1-2L	Apply at 75-150mm tall. Repeat in 10-14 days.
CHERRIES	1-2L	Apply at post harvest only.
CITRUS	1-2L	Best applied at Spring and Autumn flush. Repeat if required.
CORN/MAIZE	1-2L	Apply when knee high and repeat before tasselling.
CUCUMBER	1-4L	Apply at first bloom and repeat 0.5L after each picking.
GARLIC	1-3L	Apply when 75-175mm tall and repeat as required.
GRAPES	1-2L	Apply from first leaves up to full bloom. Post harvest.
LETTUCE	1-2L	Apply at 10 days to 2 weeks post-thinning.
ONIONS	1-4L	Apply when 75-150mm tall. Repeat as required.
NECTARINES/PEACH	0.5-1L	Apply post harvest only.
PEANUTS	1-2L	Apply 2-4 sprays from first bloom.
PEARS	0.5-1L	Apply at bud break to full bloom. Post harvest.
PISTACHIOS	1-2L	Apply at bud break to end of shoot elongation, dormant.

CROP	FOLIAR RATES L/ha	COMMENTS
POTATOES	2-5L	Apply at hooking, canopy closure and tuber bulking.
STRAWBERRIES	2-4L	Apply 2 weeks post planting and repeat 2-3 applications.
SUGARCANE	1-2L	Apply when sufficient foliage is present.
TOMATOES	1-4L	Apply from first truss and repeat as required.
WATERMELONS	0.5-1.5L	Apply up to first bloom.
ZUCCHINIS	1-3L	Apply 2 weeks post thinning and repeat 7-10 days.

Broadacre - Foliar applications per min water/hectare 50L.

CROP	FOLIAR RATES L/ha	COMMENTS
CANOLA	1-3L	Apply from 4th leaf up to pod formation.
CEREALS	2-5L	Apply mid tillering to heading.
COTTON	1-2L	Apply 6-8th leaf stage and repeat at boll formation.
RICE	3-7L	Apply 35-55 days after emergence.

**FERTIGATION:** May be applied at 7-20L/ha via overhead or dripper irrigation. Apply Zintron during middle 1/3 of irrigation cycle. Zintron can be applied with SprayCal N or 42N solutions.

ANALYSIS	AUSTRALIA W/W %	INTERNATIONAL W/W %
Nitrate N	10.5	8.0
Ammonium N	7.8	5.95
Urea N	0.7	.53
Total Nitrogen N	19	14.5
Zinc	6.6	5.0

Specific gravity +/- 0.01 pH 3.0-4.0



**STORAGE & DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE PORT ADELAIDE, SOUTH AUSTRALIA 5015  
 PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au • A.B.N. 47 007 974 496

NOT APPLICABLE TO 1000 LITRE CONTAINERS

PRODUCED AND MANUFACTURED IN AUSTRALIA

SPRAYGRO LIQUID FERTILIZERS