

24 HN-4

CAUTION



HORTICULTURE: Foliar applications per min water/hectare: 130-350 litres. Tree crops may require higher rates (500-1000L). The rates are to be adjusted according to tissue and or soil analysis results.

CROP	FOLIAR RATES L/ha	FERTIGATION RATES L/ha	COMMENTS
AVOCADOES TROPICAL FRUIT BANANAS	4 2-5	15 15	Apply at 7-14 days intervals from fruit set through to harvest. Preflowering apply up to 5L/ha, after flowering apply onto 2L/ha via foliar. Repeat if necessary. (Caution - surface of recently emerged fruit is particularly prone to marking damage from spray applications made to very rapidly growing fruit in hot conditions with direct light. Recommended to bag all emerged bunches prior to spraying to minimise risk of fruit marking).
BRASSICA'S	4	10-15	Apply at early head development to harvest every 14 days.
CUCRITS GARLIC ONIONS	2-3 3-4	5-10 12-15	Apply at fruit set onwards every 7-14 days til harvest. Apply from when sufficient leaf area exists to intercept spray and at bulb development through to harvest.
LEAFY GREENS OLIVE TREES PAW PAW	3-5 2-3	8-12 2-20 8-12	Apply every 7-14 days from mid-crop to harvest. Apply from early through to late fruiting. Apply at end of flowering and 21 days after flowering as required.
PEANUTS	n/a	10-20	Apply at planting.

CROP	FOLIAR RATES L/ha	FERTIGATION RATES L/ha	COMMENTS
PINEAPPLES	2-3	15-10	Apply at 14-21 day intervals from fruit set through to harvest.
POTATOES	2-3	4-8	Apply every 7-14 days from tuber initiation to harvest.
GRAPES	3	12	Apply from pea size berries, colouring up to 1 month before harvest.
STRAWBERRIES TOMATOES	2-3 2-3	10-15 10-15	Apply every 7-10 days from flower to harvest. Apply every 7-10 days from 2nd truss formation to harvest.

BROADACRE: Rate of foliar application per min water/hectare: 50 litres.

COTTON @ 6-8 litres/hectare at flowering and boll set. Repeat as necessary through to maturation.

LUCERNE @ 4-5 litres/hectare when sufficient leaf area exists to spray, prior flowering and after hay cutting/grazing.

LEGUME GRAINS @ 4 litres/hectare during pod development.

TURF @ 400-500mls/100m². Water dilution 1:100.

ANALYSIS	w/v%
Nitrogen (N) as urea lo bi	5
Phosphorus (P) as phosphate	10
Potassium (K) as potassium phosphate	27

ANALYSIS	w/v%
Iron (Fe) as EDTA	0.02
Zinc (Zn) as EDTA	0.018
Manganese (Mn) as EDTA	0.02
Copper (Cu) as EDTA	0.019
Boron (B) as Borate	0.017
Molybdenum (Mo) as Molybdate	0.002

Specific Gravity 1.44 ± 0.01

Impurities
 Se < 0.001%
 Cd < 1mg/kg
 Pb < 2mg/kg
 Hg < 0.2mg/kg
 F < 40g/kg P
 Biuret <15g/kg N



NOT APPLICABLE 1000 LITRE CONTAINERS

Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

Not classified as a hazardous substance according to the criteria of Safe Work Australia.
Prevention: Wear protective gloves/protective clothing/eye protection/face protection.
Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

SPRAYGRO LIQUID FERTILIZERS
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5015 • PO BOX 494
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au
PRODUCED AND MANUFACTURED IN AUSTRALIA



DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

Batch No.