



SPRAYGRO LIQUID FERTILIZERS  
 52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE PORT ADELAIDE, SOUTH AUSTRALIA 5015  
 PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@plicknowl.com.au • www.spraygro.com.au • A.B.N. 47 007 974 496

PRODUCED AND MANUFACTURED IN AUSTRALIA

Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

**STORAGE & DISPOSAL:**

| ANALYSIS OF THE KELP USED IN THE PRODUCTION OF SPRAYGRO BULL KELP (DURVILLERA POTORUM) |  |
|--|--|
| <b>AMINO ACIDS</b>   | Ascorbic Acid<br>B1 Vitamin<br>B2 Vitamin<br>E Vitamin<br>Nicotin<br>Cholin<br>Iodine<br>Alginates   |
| <b>TRACE ELEMENTS</b>  | Zinc<br>Magnesium<br>Copper<br>Manganese<br>Iron<br>Calcium<br>Sulphur<br>Boron<br>Selenium<br>Cobalt  |
| <b>ORGANIC MATTER ON:</b>  | Total Solids ..... 76%<br>Crude Fibre ..... 4.9L 0.4%<br>Sodium Chloride ..... 7% W/V<br>Nitrogen ..... 5% W/V<br>Nitrogen as Urea ..... 1% W/V<br>14% Di Methylene as Urea<br>Phosphorus ..... 0.02% W/V<br>Potassium ..... 0.18% W/V |

| CROP        | COMMENTS   |
|-------------|--|
| POTATOES    | Apply at 2-3L/ha. Apply when sufficient canopy exists to wet/cover foliage with spray. |
| GRAPE       | Apply at 4-6L/ha. Apply before flowering, then at regular intervals.                   |
| CITRUS      | Apply at 4-6L/ha. Apply just before bloom then at 3-4 week intervals.                  |
| OLIVES      | Apply at 5L/ha. Apply before bloom and repeat before fruit set.                        |
| ORNAMENTALS | Apply at 400ml per 100 litres of spray. Apply weekly as required.                      |
| FERTIGATION | Apply at 10-50L/ha.  |
| CROP        | COMMENTS   |
| TURF        | Apply at 60-100mls/100m <sup>2</sup> water dilution 1:100.                             |

| CROP   | COMMENTS   |
|--|--|
| BROADACRE: Rate of application per min water/hectare: 50 litres.     |  |
| CEREALS  | Apply at 5L/ha. Apply prior to flowering for optimum benefit.  |
| LUCERNE  | Apply at 4-6L/ha. Apply May/June as soon as sufficient leaf area exists to spray.  |
| COTTON   | Apply at 3L/ha at seeding and 4-6 leaf stage.  |
| HORTICULTURE: Rate of application per min water/hectare: 350 litres. |  |
| CROP   | COMMENTS   |
| POME FRUIT/STONE FRUIT   | Apply at 2-4L/ha. Apply one spray before bloom and repeat after bloom.   |
| ALMONDS  | Apply at 2-3L/ha. Apply one spray before bloom and repeat after bloom.   |
| STRAWBERRIES   | Apply at 10L/ha via fertigation 1-2 weeks after transplanting, then foliar apply at 5L/ha at green bud and again at flowering. |
| VEGETABLES   | Apply at 2-3L/ha. Apply when sufficient leaf area is available to intercept spray.   |
| GLASSHOUSE VEGETABLES  | Apply at 300-600mls per 100 litres of spray. Apply fortnightly during active growth stage.                                     |

**LIQUID FERTILIZER**

**BULL KKELP PLUS ++**

**LIQUID KELP SYRUP - ORGANIC PLANT GROWTH REGULATOR**

○ 20 Litre NET ○ 200 Litre NET

**CAUTION**

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**Safety glasses, gloves and protective clothing should be worn when using this product.**

**DISCLAIMER/CONDITIONS OF SALE:** 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred of suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.



BATCH No.