



Suggested Nutrient Applications for:



DRIED FRUIT

Latest Update: 10/8/09

Growth Stage	Bud Burst to Flowering	Early Flowering	Flowering	Fruit Set	Berries forming pea size	Variation	Post Harvest
Product % Nutrient (N-P-K) W/V							
Smartrace Magnesium 5.6-0-0, 5 Mg		4					
Pick 15-42 0-15-42					4-6	3-4	
Smartrace Calcium 3.5-0-0, 5 Ca					3	3	
Smartrace Zinc Manganese 5 Zn, 5 Mn, 5.4 S	2 applications at 4lt/ha	4		4			
Smartrace Spraytrace 8 0.2-0-0, 1.35 Mg, 1.34 Zn, 1.34 Fe, 1 Mn, 0.5 B, 0.02 Mo, 0.17 Cu, 2.1 S							4-6
Result 10-5-7 + TE	8-10 X 2 applications			4		4	10
Nitro 35 35 – 0 – 0							15
Water Rate: 1000 litres per Hectare.							
<div style="display: flex; justify-content: space-around; align-items: center;"> = Foliar = Bandspray or Fertigation </div>							
<p>Important Note:</p> <p>(1) The product label should be consulted before use of any of the products referred to on this sheet, always refer to the label when applying SprayGro Liquid Fertilizer products.</p> <p>(2) This information sheet is a guide only as SprayGro Liquid Fertilizers neither makes, nor authorizes any agent or representative to make any other warranties, express or implied, other than those stated on the product label.</p> <p>(3) SprayGro Liquid Fertilizers shall not be liable for any results, loss, or damage whatsoever, whether consequential or otherwise through the use or application of products and/or materials referred to herein.</p> <p>(4) We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray. Combination sprays can induce crop phytotoxicity.</p> <p>(5) Apply at times of low light intensity, early morning or preferably during the evening. Avoid applications to wet leaves, stressed crops and on exceptionally hot, dry or sunny days. Overcast, calm and cool conditions are best.</p> <p>(6) E.&O.E.</p>							