

**BROADACRE:**  
 LUCERNE & PASTURE @ 5 litres/hectare in 100-250- litres water per hectare.  
 COTTON @ 2-5 litres/hectare at 2 weeks after first bloom at 14 day intervals x 3 applications. Water rate 30-100 litres/hectare.

**HORTICULTURE:**  
 APPLES & PEARS @ 3-5 litres/hectare in 500-1000 litres of water after petal fall and repeat as necessary.  
 BANANAS @ 10-12 litres/hectare. Apply after flowering in minimum of 200 litres water.  
 BRASSICAS @ 3-5 litres/hectare. Apply at 4-6 true leaves and repeat as necessary at 7-10 day intervals in 250-750 litres water per hectare.  
 CARROTS & ONIONS @ 5 litres/hectare in 500 litres of water when crop is 10-15cm high in 2-500 litres water.  
 CITRUS @ 2 litres/hectare. Apply after fruit set as necessary at 14-21 day intervals in 1000 litres water per hectare.  
 MANGO @ 2 litres/hectare. Apply after fruit set as necessary at 14-21 day intervals in 400 litres water per hectare.  
 PINAPPLES @ 5 litres/hectare. Apply 4-6 weeks before harvest in 4-600 litres water.  
 POTATOES @ 3-5 litres/hectare in 500 litres of water at tuber formation and tuber bulking at 7-10 day intervals.  
 STONE FRUITS @ 3-5 litres/hectare in 500 litres of water at fruit set and repeat applications at 10-14 day intervals, water 500-1000 litres/hectare.  
 STRAWBERRIES @ 5 litres/hectare. Apply after flowering as necessary at 7-10 day intervals in 500-1000 litres water.  
 TOMATOES @ 3-5 litres/hectare in 500 litres of water at fruit set at 7-10 day intervals in 2-500 litres water.  
 VINES @ 3-5 litres/hectare in 500 litres of water 10-14 days after flowering.  
 FERTIGATION Up to 50 litres/hectare, repeat as necessary at 10-14 day intervals.

**TURF:**  
 TURF @ 200-300mls per 100m<sup>2</sup> / water dilution 1:100. Repeat applications as required.

**NURSERY:**  
 Rate of application: Soil drench nursery stock @ 5 litres per 1000 litres water.  
 Soil drench nursery stock @ 8 litres per 1000 litres water.  
 Foliar apply @ 1.25-2 litres per 1000 litres water.



**LIQUID FOLIAR FERTILIZER**

**Hi-K 41.5**

POTASSIUM AS POTASSIUM CARBONATE

**N P K**  
**0 : 0 : 41.5% W/W**

20 Litre  200 Litre  1000 Litre

**CAUTION Hi-K 41.5** is a high alkaline product with high ph of 11.

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- **Compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **Hi-K 41.5 can not** be mixed with calcium or magnesium materials.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**Safety glasses, gloves and protective clothing should be worn when using this product.**

**CAUTION: KEEP AWAY FROM CHILDREN.** Wear goggles or a face mask when using **Hi-K 41.5**. Avoid contact with skin. If swallowed give water, induce vomiting and seek medical attention.

**CONDITIONS OF SALE:** 1) Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than the officer of the Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2) Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

**BATCH No.**

SPRAYGRO LIQUID FERTILIZERS PTY. LTD.  
 52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE  
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496  
 PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@picknowl.com.au

**PRODUCED AND MANUFACTURED IN AUSTRALIA**