

BATCH No.



9 533056 006032

# MAXI-PHOS 16

N P K  
10 : 16 : 0 w/v

+ chelated trace elements

20 Litre NET  200 Litre NET  1000 Litre NET

## LIQUID FERTILIZER

# SPRAYGRO<sup>®</sup>

30 HN-3

### CAUTION

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**Safety glasses, gloves and protective clothing should be worn when using this product.**

**DISCLAIMER/CONDITIONS OF SALE: 1)** Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and/or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly, in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. **2)** The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. **3)** The customer understands that any representation of the type set out above is, on the basis that the customer must independently verify the accuracy thereof. **4)** The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. **5)** Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. **6)** Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.

#### BROADACRE: Rate of application per min water/hectare: 50 litres.

- CEREALS @ 4-6 litres/hectare. Apply as a nutrient boost 6 weeks after emergence.
- LUCERNE @ 4-6 litres/hectare. Apply May/June as soon as sufficient leaf area exists to spray.
- COTTON @ 5-20 litres/hectare as a seedling starter. Apply 2-7 litres/hectare 4-6 leaf stage.
- PASTURE @ 4-6 litres/hectare

#### SEED DRESSING: Rate of application.

- SEED @ 8 litres/tonne of seed.

#### HORTICULTURE: Rate of application per min water/hectare: 350 litres.

- POME FRUIT @ 2-4 litres/hectare. Apply one spray before bloom, repeat after bloom as necessary.
- STONE FRUIT @ 2-3 litres/hectare. Apply when deficiency exists as necessary.
- ALMONDS @ 2-4 litres/hectare. Apply one week after transplanting then again prior to bloom if required.
- STRAWBERRIES @ 2-4 litres/hectare. Repeat at regular intervals during growing season. Use 3-5 applications in conjunction with normal spraying program.
- VEGETABLES @ 2-4 litres/hectare. Apply when sufficient canopy exists to wet/cover foliage with spray.
- POTATOES @ 2-4 litres/hectare. Apply at least once before flowering, then at regular intervals.
- GRAPES @ 4-6 litres/hectare. Apply just before blossom, at petal fall and or more times at 4 weekly intervals.
- CITRUS @ 3-4 litres/hectare. Apply as necessary when deficiency exists.
- OLIVES @ 10-50 litres/hectare.
- FERTIGATION
- TURF

#### TURF: Rate of application:

- TURF @ 400-500mls/100m<sup>2</sup> water dilution 1:100.



#### STORAGE & DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

#### SPRAYGRO LIQUID FERTILIZERS

52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE  
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496  
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au

NOT APPLICABLE TO 1000 LITRE CONTAINERS

**PRODUCED AND MANUFACTURED IN AUSTRALIA**

**Application recommendations for Maxi-phos 16:**  
**apply with sufficient water for thorough coverage of leaves.**