



BATCH NO.

**Safety classes, gloves and protective clothing should be worn when using this product.** (1) Foliar fertilizers are not to be used on plants that are under stress or damage including (but not limited to) frost damage, drought stress, insect damage, or other damage. (2) The customer warrants that there has been no reliance by the customer upon any representation made by SprayGro or upon and descriptions, illustrations or specifications contained in SprayGro's Product Profile or any other document produced by SprayGro. (3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. (4) The customer shall at all times indemnify and hold harmless SprayGro from and against all loss (including reasonable legal costs and expenses) or liability (reasonably incurred or suffered) by SprayGro arising from any proceedings or (including reasonable legal costs and expenses) or liability (reasonably incurred or suffered) by SprayGro arising from any conditions of use. No one other than an authorized officer of SprayGro is authorized to make any warranty, guarantee or direction concerning this product. (5) Before using please obtain a current Product Profile which is available from SprayGro or our Distributor.

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture.
- Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket) and check for precipitation (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**CAUTION**



**LIQUID FERTILIZER**

**MOLYPHOS**

**MOLYBDENUM IS ESSENTIAL FOR NITROGEN ASSIMILATION.**

**GUARANTEED MINIMUM ANALYSIS**  
 MOLYBDENUM.....25.0% W/W.  
 PHOSPHORUS.....11.0% W/W.

**5 Litre NET**  **20 Litre NET**  **200 Litre NET**

MolyPhos application recommendations: Apply to the following crops where a deficiency exists. **MINIMUM DILUTION:** A dilution of 1:100 means 1 part product : 100 parts water. In hot weather use a higher dilution rate where applicable.

CROP	FOLIAR RATES ml/ha	MINIMUM DILUTION	COMMENTS
APPLES	250-350	1:1000	For best results apply onto new foliage.
ASPARAGUS	125-250	1:700	Apply when required.
AVOCADOS	250-500	1:1000	Apply when required or apply at 4-5 leaf stage.
BRASSICAS	250-500	1:400	Apply at 4-6 leaf stage. Follow up applications at 7-10 day intervals as required.
CAPSAICUM	125-250	1:800	Apply when required or apply at 4-5 leaf stage.
CELERY	80-200	1:1000	Apply when required or apply at 4-5 leaf stage.
CITRUS	150-250	1:2000	For best results apply onto new foliage when required.
COTTONCEREALS	200-300	1:500	For best results apply onto new foliage when required.
CUCURBITS	250-500	1:400	Apply when required or apply at 4-5 leaf stage.
GRAPE VINES	200-350	1:1500	For best results apply onto new foliage when symptoms are exposed.
LEGUMES	250-500	1:400	Apply when early growth is approx. 150mm tall to ensure effective movement into nitrogen fixing nodules.
LETTUCE	250-500	1:400	Apply at 2-4 leaf stage. Follow up applications at 7-10 day intervals as required.
NUT CROPS	100-250	1:2000	For best results apply onto new foliage when symptoms are exposed.

CROP	FOLIAR RATES ml/ha	MINIMUM DILUTION	COMMENTS
OLIVES	80-250	1:2000	For best results apply onto new foliage when required.
*PASTURE/ LUCERNE	150-250	1:800	Apply when early growth is approx. 150mm tall to ensure effective movement into nitrogen fixing nodules. Apply at least 2 weeks before cutting.
POTATOES	100-250	1:1000	Apply as symptoms are exposed or apply at 4-5 leaf stage.
STONE FRUIT	50-200	1:2000	For best results apply onto new foliage when required.
STRAWBERRIES	125-250	1:400	Apply when required or apply at 4-5 leaf stage.
TOMATOES	200	1:250	Apply when required or apply at 4-5 leaf stage.
TREE CROPS	250-500	1:1000	For best results apply onto new foliage when required.
VEGETABLES	125-250	1:400	Apply when required or apply at 4-5 leaf stage.
ZUCCHINI	100-250	1:400	Apply when required or apply at 4-5 leaf stage.

ANALYSIS	AUSTRALIA W/W %	INTERNATIONAL W/W %
Molybdenum	25.0	16.0
Phosphorus	11.0	(P <sub>2</sub> O <sub>5</sub> ) 16.0

Specific gravity 1.57 pH 5.0-6.0

**CAUTION:**  
 This product is harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist. Warning: May cause corrosion of galvanised spray equipment. Thorough washing of equipment after use is recommended. **\*Warning: Excessive use of molybdenum can be harmful to stock. Plant levels of molybdenum can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period. Check rate and frequency of molybdenum use with appropriate authorities.**

**STORAGE & DISPOSAL:**  
 Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

**SPRAYGRO LIQUID FERTILIZERS**  
 52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE PORT ADELAIDE, SOUTH AUSTRALIA 5015  
 PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au • A.B.N. 47 007 974 496

**PRODUCED AND MANUFACTURED IN AUSTRALIA**



NOT APPLICABLE TO 1000 LITRE CONTAINERS

271RS-1