

SMART RELEASE

NITROGEN™
SERIES

NITRO-11 HI K

NPK **11-0-15** W/V

Contains 85% Smart Release Nitrogen™ from Methylene Urea

Low salt index formulation ■ Controlled release ■ Non leaching
CONTAINS 12% SULFUR



NOT APPLICABLE TO 1000 LITRE CONTAINERS

20 Litre NET 200 Litre NET 1000 Litre NET

CAUTION

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture.
- Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

Safety glasses, gloves and protective clothing should be worn when using this product.

DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro or by involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.



BATCH No.

CROP	FOLIAR FERTIGATION RATES L/ha	COMMENTS
ALMONDS	5-15L	Apply at first full leaf and as needed during growing season.
APPLES	5-15L	Apply at first full leaf and as needed during growing season.
AUSTRALIAN NATIVES	3-4L	Apply mid to late vegetative. Repeat at flowering.
CAPSICUMS	4-6L	Apply at 7-14 days from fruitset onwards.
CITRUS	5-15L	Apply at first full leaf and as needed during growing season.
CUCUMBERS	4-6L	Apply at 7-14 days from fruitset onwards.
CUCURBITS	2-3L	Apply every 7-14 days from fruitset onwards.
MAIZE	5-6L	Apply prior to tasselling. Repeat in 10-14 days.
ONIONS	3-4L	Apply from when sufficient leaf area exists to intercept spray and from bulb development onwards.
POTATOES	4-6L	Apply 14-30 days from tuber initiation to harvest.
STRAWBERRIES	4-6L	Apply every 14-21 days from flower to harvest.
TOMATOES	4-6L	Apply every 7-10 days from 2nd truss formation.
VEGETABLES & ROW CROPS	3-4L	Apply from 3rd to 4th leaf stage every 10-14 days based on crop requirements.

HORTICULTURE: Foliar applications per min water/hectare generally required: 150-300L. Tree crops may require higher rates (500-1000L).

Application rates for Nitro-11 Hi K
CAUTION: Do not apply to foliage of crops sensitive to (foliar burn) to sulphur.
Do not apply with oil sprays.

CROP	COMMENTS
COTTON	Apply 6-8L/ha at flowering and boll set. Repeat as necessary through to maturation.
LUCCERNE	Apply 10-14L/ha when sufficient leaf area exists to spray prior to flowering and after hay cutting/grazing.
LEGUME GRAIN	Apply 4L/ha during pod development.

BROADACRE: Foliar rate of application per min water/hectare: 80 litres.

CROP	COMMENTS
TURF	Low feed: Apply 90L/ha apply min 1:25 water dilution. Medium feed: Apply 130L/ha apply min 1:25 water dilution. High feed: Apply 180L/ha apply min 1:25 water dilution.

TURF: Can be applied with low water rates as a foliar or use higher water rates to drench product in the soil. Smart Release Nitrogen enhances the uptake of other nutrients. Apply the following rates at 4-6 week intervals for optimum turf growth.

ANALYSIS	AUSTRALIA W/V %	INTERNATIONAL W/V %
Nitrogen (N)	11	8.4
Phosphorous (P)	15	10
Potassium (K)	0	14
Sulphur (S)	12	10

Specific gravity 1.31 pH 9-10

Should you require a higher Nitrogen amount, tank mix with Smart Release Nitrogen™ Nitro-37.

PRODUCED AND MANUFACTURED IN AUSTRALIA

SPRAYGRO LIQUID FERTILIZERS
52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE, PORT ADELAIDE, SOUTH AUSTRALIA 5015
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@picknroll.com.au • www.spraygro.com.au • A.B.N. 47 007 974 496

Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

STORAGE & DISPOSAL: