



Suggested Nutrient Applications for:

Potatoes assisted with Granular Planting & Sidedress

Latest Update: 23/03/06

Growth Stage	Planting	Germination	Hilling	Hooking	Tuber Initiation	Full Canopy	Bulking	Desiccation	Notes
Product % Nutrient (N-P-K) W/V									
Pick 20:40 0-20-40 + 0.5 Fulvic Acid		20	20	20					High analysis P & K in highly plant available form to drive early root growth
K Blast 30 0-0-30, S 24		Rate			30	30	30		High analysis potassium with good uptake and low salt index
Firmrite SprayCal Mag 12-0-0, Ca 14, Mg 2.5			Litres/Hectare	50	50	50			Calcium Nitrate with very high levels of Magnesium
Boron 15 6-0-0, B 15						1.5		2	Non-sodium form of Boron, which drives Calcium uptake, tuber set, bulking & quality.
Spray Cat 8 0-0-0, Ca 8, Su 14			0.5					Calcium Thiosulphate Calcium, highly available for better uptake and transport.	
Gripper Trace 5 1-0-0, S 6.0, Mg 2.0, Zn 5.0, Mn 3.0, Cu 1.0, Fe 1.0			5		5		8	EDTA Chelated multi-trace elements product in an ideal ratio.	

* = Product to Product Compatibility when diluted with water at correct rates

Minimum Water Rate: Foliar Application 1:50 Fertigation 1:100 = Foliar (Lt/Ha) = Fertigation (Lt/Ha)

Important Note:

- (1) The product label should be consulted before use of any of the products referred to on this sheet, always refer to the label when applying SprayGro Liquid Fertilizer products.
- (2) This information sheet is a guide only as SprayGro Liquid Fertilizers neither makes, nor authorizes any agent or representative to make any other warranties, express or implied, other than those stated on the product label.
- (3) SprayGro Liquid Fertilizers shall not be liable for any results, loss, or damage whatsoever, whether consequential or otherwise through the use or application of products and/or materials referred to herein.
- (4) We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray. Combination sprays can induce crop phytotoxicity.
- (5) Apply at times of low light intensity, early morning or preferably during the evening. Avoid applications to wet leaves, stressed crops and on exceptionally hot, dry or sunny days. Overcast, calm and cool conditions are best.
- (6) E.&O.E.