

Rate of application: TURF @ 500mls/100'water dilution 1:100.

**TURF:**  
OLIVES  
STRAWBERRIES @ 2 litres/hectare. Apply a week after transplanting, then prior to bloom if necessary.  
APPLES, PEARS @ 4-8 litres/hectare. Apply 3-4 weeks after full bloom then repeat as necessary.

Rate of application per min water/hectare: 350 litres  
STONE FRUIT, NUT TREES.

**FRUIT CROPS:**

CITRUS @ 2-4 litres/hectare. Apply to thoroughly cover foliage as a split application a fortnight prior to bloom, then again after bloom if necessary.  
Repeat if necessary.  
GRAPES @ 2-4 litres/hectare. Apply as thorough cover spray a fortnight to three weeks before bloom. Do not apply during bloom.  
BEANS @ 2-3 litres/hectare. Apply when the plant is between second and third leaf stage. Repeat as necessary.  
CUCUMBERS, MELONS @ 2-3 litres/hectare. Apply during the flowering phase when fruit is being set.  
CELERY @ 2-3 litres/hectare. Apply one week to a fortnight after emergence for seeded crops. Repeat as necessary.  
CARROTS, LETTUCE, applications. Repeat as necessary.

POTATOES @ 1-2 litres/hectare. Apply within a month of emergence and before flowering. Split applications are preferred over single ONIONS @ 2-3 litres/hectare. Apply when plant is 7-15cm high prior to blossoming, then again as necessary when fruit has set. VEGETABLES, TOMATOES @ 2-3 litres/hectare. Apply in initial growing stages when deficiency symptoms occur and repeat as necessary.

Rate of application per min water/hectare: 300 litres

**HORTICULTURE:**

LUCERNE, PASTURES @ 2-3 litres/hectare. Apply in early spring when rapid growth period begins. Repeat sprays at least once per cutting early in growth cycle of each cutting. Do not apply at bloom.  
CEREAL, COTTON @ 1-3 litres/hectare. Apply to actively growing plants when sufficient leaf area exists to spray. Split applications are preferred to single applications.

Rate of application per min water/hectare: 50 litres

**BROADACRE:**

Apply with sufficient water for thorough coverage of leaves. APPLICATION RECOMMENDATIONS: SPRAYGRO SMARTTRACE CALCIUM is suitable for the following crops where a calcium deficiency exists.



**CAUTION**

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- **Compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**Safety glasses, gloves and protective clothing should be worn when using this product.**

**CONDITIONS OF SALE:** 1) Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than the officer of the Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2) Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

SPRAYGRO LIQUID FERTILIZERS PTY. LTD.

52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE  
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496  
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@picknowl.com.au

BATCH No.

PRODUCED AND MANUFACTURED IN AUSTRALIA