



BATCH No.

DISCLAIMER/CONDITIONS OF SALE: (1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. (2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. (3) The customer understands that any representation of the type set out above is on the basis that the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised office of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. (6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.

Safety glasses, gloves and protective clothing should be worn when using this product.

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

- **DO NOT** apply during the heat of the day.
 - **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
 - **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
 - **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
 - Use the correct label spray rates of any pesticide used with Spraygro product.
 - Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
 - bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
 - **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
 - Use Spraygro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. In damage to the plant.
- The following conditions must be observed in order to apply Spraygro product blends successfully. Failure to follow these instructions may result

CAUTION



SMARTTRACE
CHELATED ELEMENTS

ZINC

CONTAINS
4.9% W/W
SULPHUR

GUARANTEED MINIMUM ANALYSIS
ZINC AS NATURAL PHENOLIC COMPLEX 10% W/W

20 Litre NET ● 200 Litre NET ● 1000 Litre NET ●

APPLICATION RECOMMENDATIONS: SPRAYGRO SMARTTRACE ZINC. Apply when a Zinc deficiency exists, with sufficient water for thorough coverage of leaves.

BROADACRE:

- Rate of application per min water/hectare:** 50 litres
- PASTURES @ 2-3 litres/hectare.
 - CEREALS GRAINS @ 2-3 litres/hectare.
 - LEGUME GRAINS @ 2-3 litres/hectare.
 - COTTON @ 2-3 litres/hectare.

TURF:

- Rate of application per min water/hectare:** 300 litres
- TURF @ 150mls/100m²/dilution 1:100/

SEED DRESSINGS: Suggested application rate:

- WHEAT, BARLEY, OATS @ 5 litres per tonne of seed.
- PEAS, LUPINS, BEANS @ 5-6 litres per tonne of seed.
- MAIZE, RICE @ 6-8 litres per tonne of seed.

HORTICULTURE:

- Rate of application per min water/hectare:** 300 litres
- POTATOES @ 2-4 litres/hectare.
 - CITRUS @ 2-4 litres/hectare. Apply during Spring/Autumn flush as required.
 - ALMONDS @ 2-3 litres/hectare. Apply at petal fall, then twice at 4 week intervals.
 - Avoid spraying on Challiston Variety.*
 - GRAPES @ 3-5 litres/hectare. Apply as through cover spray 2-3 weeks before bloom. Do not apply during bloom. Repeat as necessary.
 - STRAWBERRIES @ 1-2 litres/hectare after transplanting.
 - VEGETABLES @ 2-4 litres/hectare.
 - BANANAS @ 3-4 litres/hectare early canopy and 1-2 litres/hectare after flowering. (Caution - surface of recently emerged fruit is particularly prone to marking damage from spray applications made to very rapidly growing fruit in hot conditions with direct sunlight. Recommended to bag all emerged branches prior to spraying to minimise risk of fruit marking).
 - AVOCADOS @ 2-3 litres/hectare spring flush and again after harvest.

ANALYSIS	AUSTRALIA W/W %	INTERNATIONAL W/W %
Zinc Zn	10	7.9

Specific gravity 1.27 pH 3.4 - 4.5

STORAGE & DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information. **The rates must be adjusted according to tissue and or soil analysis results.**

SPRAYGRO LIQUID FERTILIZERS

76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA



NOT APPLICABLE TO 1000 LITRE CONTAINERS

09 SR-7