

Latest Update: 22/05/07



Suggested Nutrient Applications for:

Table Grapes Red

Growth Stage	Pre Bud Burst	Bud Burst (BB)	4w Post BB	6w Post BB	Pre-flowering	Flowering	Fruit Set	Pea Size Berries	Bunch Closure	Veraison	Pre-Harvest	Post-Harvest	Notes
Product	% Nutrient (N-P-K) w/v												
Humic Acid 12 12 Ha	12												Ameliorates soil structure and properties to retain nutrients and moisture
MaxiPhos Injecta 23 16-23-0	15				15							15	Ammonium polyphosphate (APP) starter promotes early crop establishment. Better P delivery. Replace less efficient MAP and SSP
Result Plus 14-8-10 + 0.1 S, 0.01 Mg, 0.02 Fe, 0.050 Mn, 0.0175 Zn, 0.017 B, 0.017 Cu, 0.0016 Mo, 0.0017 Co			30		30		30		30			10	NPK Fertigation product to improve root growth and shoot development; fortified with EDTA trace elements
SMARTRACE® Zinc/Mang. 5Zn, 5Mn 5.4 S	3												Improves uniform budburst and growth
Smartrace Magnesium 5.6-0-0, 5Mg			3		3		3	3		3			Magnesium improves sugar translocation and chlorophyll.
SMARTRACE® Calcium 2-0-0, 5Ca							5	5					Lignosulfonate chelated calcium to improve colour and firmness.
K Blast 36 0-0-36, 8.8S		= PRODUCT TO PRODUCT COMPATIBILITY*				25	25		30				The high K fertilizer to promote colour and fruit development. Potassium increase sugar content.
Firmrite SprayCal Mag 12-0-0, 14Ca, 2.5Mg				25				25		20			Calcium and magnesium in ideal ratio to improve berry size, strength and resistance to pest infestation.
SMARTRACE® Spraytrace 8 1 Mn, 1.34, Zn, 0.17 Cu, 1.35 Mg, 0.5 B, 0.05 Fe, 1.34, 0.02 Mo, 2.1S, 0.2N				5		5			5	5		5	Complete micronutrient fertilizer to remove deficiencies and maximize growth.

**Rate
Litres/Hectare**

* = Product to Product Compatibility when first diluted with water at correct label rate
 Minimum Water Rate: Foliar Application 1:50 Fertigation 1:100 = Foliar (L/ha) = Fertigation (L/ha)

- (1) The product label should be consulted before use of any of the products referred to on this sheet, always refer to the label when applying SprayGro Liquid Fertilizer products.
- (2) This information sheet is a guide only as SprayGro Liquid Fertilizers neither makes, nor authorizes any agent or representative to make any other warranties, express or implied, other than those stated on the product label.
- (3) SprayGro Liquid Fertilizers shall not be liable for any results, loss, or damage whatsoever, whether consequential or otherwise through the use or application of products and/or materials referred to herein.
- (4) We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray. Combination sprays can induce crop phytotoxicity.
- (5) Apply at times of low light intensity, early morning or preferably during the evening. Avoid applications to wet leaves, stressed crops and on exceptionally hot, dry or sunny days. Overcast, calm and cool conditions are best.
- (6) E.&O.E.