

50 HN-2

CAUTION

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**Safety glasses, gloves and protective clothing should be worn when using this product.**

**DISCLAIMER/CONDITIONS OF SALE:** 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.

9 333056 002591



BATCH No.

CROP	FOLIAR RATES L/ha	FERTIGATION RATES L/ha	COMMENTS
LETTUCE	2-3L	8-10L	Apply from emergence, continue 14 day intervals until harvest.
NUT CROPS	4-6L	12-20L	Apply 2 weeks post bloom, repeat at regular intervals during nut development. Repeat post-harvest.
OLIVES	5-7L	10-20L	Apply 2 weeks prior to fruit set, at fruit set and further applications as required.
PEARS	2-3L	7-10L	Apply at petal fall. Continue every 14 days to harvest.
POTATOES	5-7L	20-40L	Apply 3 weeks after emergence. Repeat at hooking, canopy closure and bulking.
STONE FRUITS	3-4L	10-15L	Apply post-bloom. Continue from fruit set at regular intervals.
STRAWBERRIES	2-4L	10-20L	Apply 2 applications prior to flowering with further applications at regular intervals.
TOMATOES	3-4L	10-20L	Apply every 14-21 days from 2nd truss to harvest.
TROPICAL FRUIT	2-4L	10-20L	Spray before bloom. Repeat at fruit set until 7 days before harvest.
VEGETABLES	2-3L	7-10L	Apply 3 weeks after emergence or post-transplant. Repeat at regular intervals.
ZUCCHINI	2-3L	7-12L	Two applications prior to flowering. Repeat during fruit elongation until 1 week before harvest.

**TURF - Rate and time of application:**  
500-700mls/100m<sup>2</sup>/water dilution 1:100. Repeat applications as required.

ANALYSIS	AUSTRALIA W/W %	INTERNATIONAL W/W %
Nitrogen (N)	0	0
Phosphorous (P)	0	0
Potassium (K)	36	(K2O) 30

Specific gravity 1.42 + - 0.010

STORAGE & DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

SPRAYGRO LIQUID FERTILIZERS

76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE PORT ADELAIDE, SOUTH AUSTRALIA 5015  
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au • A.B.N. 47 007 974 496

PRODUCED AND MANUFACTURED IN AUSTRALIA

LIQUID FERTILIZER

IK BLAST 36

POTASSIUM AS CITRATE, THIOSULFATE & CARBONATE

N P K

0 : 0 : 36 + 8.8 S w/v

20 Litre NET 200 Litre NET 1000 Litre NET

**HORTICULTURE:** Foliar applications per min water/hectare generally required: 150-300L. Tree crops may require higher rates (500-1000L).

CROP	FOLIAR RATES L/ha	FERTIGATION RATES L/ha	COMMENTS
ALMONDS	5-7L	15-30L	Apply 2 weeks post-bloom. Repeat 2-3 times every 4 weeks.
APPLES	5-7L	15-25L	Apply at petal fall. Continue at 7-14 day intervals ntil harvest.
ASPARAGUS	4-6L	8-12L	Apply 7 days from 1st fern. Repeat at spear emergence.
AVOCADOES	3-5L	10-15L	Apply 3 weeks prior to flowering. Repeat every 3 weeks from fruit set to harvest.
CARROTS & ONIONS	3-4L	10-12L	Apply weekly from root differentiation until 1 week before harvesting carrots. Apply from bulb development at regular fortnightly intervals in onions.
BROCCOLI	2-3L	7-15L	Apply every 2 weeks until head formation.
CAPSICUM	2-3L	9-15L	Apply every 2 weeks until flowering. Repeat at regular intervals during fruiting.
CELERY	2-3L	7-12L	Apply every 2 weeks. Stop application 1 week before harvest.
CITRUS	5-7L	10-25L	Apply prior to bloom, at petal fall and later as required.
COTTON	5-6L	10-20L	Apply prior to 1st flowering. Repeat at boll formation.
CUCUMBERS & CUCURBITS	2-3L	8-12L	Apply prior to flowering with further applications at regular intervals Do not apply a week before harvest.
GRAPE VINES	2-3L	10-25L	Apply prior to flowering with regular applications until 2-3 weeks before harvesting.

**The rates are to be adjusted according to tissue and or soil analysis results.**



NOT APPLICABLE TO 1000 LITRE CONTAINERS