


BATCH No.



DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spray Gro Liquid Fertilizer and its servants and agents ("Spray Gro") shall be under no liability to the purchaser and/or user of Spray Gro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spray Gro Product Profile or not. **2)** The customer warrants that there has been no reliance by the customer upon any representation made by Spray Gro or upon and descriptions, illustrations or specifications contained in Spray Gro's Product Profile or any other document produced by Spray Gro or upon and descriptions, illustrations or specifications contained in Spray Gro's Product Profile or that the customer must independently verify the accuracy thereof. **4)** The customer shall at all times indemnify and hold harmless Spray Gro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred of suffered by Spray Gro arising from any proceedings or threatened proceedings against Spray Gro by or involving the customer. **5)** Spray Gro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spray Gro is authorised to make any warranty, guarantee or direction concerning this product. **6)** Before using please obtain a current Product Profile which is available from Spray Gro or our Distributor.

Safety glasses, gloves and protective clothing should be worn when using this product.

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

DO NOT apply during the heat of the day.

DO NOT apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

DO NOT spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

DO NOT use when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

DO NOT apply when crop is under heavy stress from pests, heat or inadequate moisture levels.

SMARTTRACE[®]

CHELATED ELEMENT

MAGNESIUM

GUARANTEED MINIMUM ANALYSIS
MAGNESIUM AS NATURAL PHENOLIC COMPLEX 5.0% W/W

20 Litre NET ■ 200 Litre NET ■ 1000 Litre NET ■

APPLICATION RECOMMENDATIONS: Spray Gro SMARTTRACE MAGNESIUM is suitable for the following crops where a magnesium deficiency exists. Apply with sufficient water for thorough coverage of leaves.

BROADACRE: Rate of application per min water/hectare: 50 litres/ha

PASTURE @ 1-2 litres/hectare. Apply when deficiency is evident.

CEREAL GRAINS @ 3-4 litres/hectare. Apply to actively growing plants when sufficient leaf area exists to spray. Repeat as necessary. Apply mid-late tillering.

COTTON @ 3-4 litres/hectare. Apply when deficiency is evident or earlier.

LUCERNE @ 3-4 litres/hectare. Apply pre-flowering or earlier if a deficiency is evident.

CANOLA @ 3-4 litres/hectare. Apply at 4-6 leaf stage.

LEGUME CROPS @ 2-3 litres/hectare. Apply pre-flowering or earlier if a deficiency is evident.

HORTICULTURE: Rate of application per min water/hectare: 300 litres

VEGETABLES, TOMATOES @ 2-3 litres/hectare. Apply in initial growing stages when deficiency symptoms occurs.

For severe deficiencies additional applications can be made at fortnightly intervals.

CITRUS, FRUIT TREES

STONE FRUIT,

@ 2-4 litres/hectare. Apply fortnightly in springtime as a thorough spray, up until bloom and again after bloom if necessary. Repeat applications may be necessary for a severe deficiency. Best absorption occurs 1/2 – 1/3 through a flush of growth. Spray should not be applied during blossoming or harvest.

NUT TREES @ 2-4 litres/hectare. Apply 7-14 days after petal fall.

GRAPEVINES @ 2-4 litres/hectare. Apply before and after flowering.

BERRIES @ 2-4 litres/hectare. Apply at the first sign of leaf purpling symptoms, then regular sprays at 4 week intervals as required.

TURF: Rate of application:

TURF @ 100-400mls/100m²/water dilution 1:100.

NURSERY: Rate of application:

SHRUBS, ORNAMENTALS Foliar spray at 1-2 litres per 1000 litres water.

Soil application drench at 4-8 litres per 1000 litres water.

ANALYSIS	AUSTRALIA W/W %	INTERNATIONAL W/W %
Magnesium Mg	5.0	4.06
Specific gravity 1.23 pH 6.4-7.1		

STORAGE & DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

SPRAYGRO LIQUID FERTILIZERS

76 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE PORT ADELAIDE, SOUTH AUSTRALIA 5015
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • A.B.N. 47 007 974 496



NOT APPLICABLE TO 1000 LITRE CONTAINERS

PRODUCED AND MANUFACTURED IN AUSTRALIA